

Bibliografia

- ABRAMS M., HOOK S., RAMACHANDRAN B., 2002. *ASTER User's Handbook. Version 2*. Jet URL: <http://asterweb.jpl.nasa.gov>.
- ADAMCZEWSKA A.M., MORRIS S., 1998. The functioning of the Haemocyanin of the terrestrial Christmas island red crab *Gecaroidea natalis* and the roles for organic modulators. *Journal of Experimental Biology*, 201: 3233-3244.
- AGRICONSULTING, 2008. Bozza di piano di gestione del SIC/ZPS IT3320037 Laguna di Marano e Grado. Terzo rapporto intermedio. Relazione ad uso della Regione Autonoma Friuli Venezia Giulia.
- ALTOBELLI A., HUBINA T., BRESSAN E., DE SIMON G., 2007. Analisi della morfologia del sistema lagunare di Grado attraverso immagini multispettrali ASTER, *Rivista italiana di Telerilevamento*, 39: 67-75.
- ALTOBELLI A., HUBINA T., SPONZA S., SISTO A., 2008. Effect of abiotic and biotic factors on the abundance of waterbirds in Grado-Marano Lagoon (Italy). *Proc. SPIE*, Volume 7104, Wetlands and Lake Habitats.
- ANGER K., SPIVAK E., LUPPI T., 1998. Effects of reduced salinities on development and bioenergetics of early larva shore crab, *Carcinus maenas*. *Journal of Experimental Marine Biology and Ecology*, 220: 287-304.
- AOAC, 1990. Standard Methods 15th Ed.
- ARANDA X. G., MCPHEETERS R., IGOE F., MORAN B., EISENBERG A. J., 2004. Alkaline Extraction of DNA from FTA® Paper Spotted with Buccal Epithelial Cells and Whole Blood. Poster: Promega 15th International Symposium on Human ID, Phoenix (AZ).
- ARCHAWARANON M., 2004. Rapid sexing hill mynah *Gracula religiosa* by sex chromosomes. *Biotechnology* 3: 160-164.
- ARPA-FVG, 2008. L'alto Adriatico, il Golfo di Trieste, il litorale del Friuli Venezia Giulia. <http://www.arpa.fvg.it/index.php?id=140> (Query date: 21/08/2008).
- BACCETTI N., CHERUBINI G., MAGNANI A., SERRA L., 1995. Homing performance of adult and immature Dunlins *Calidris alpina* (Aves Scolopacidae) displaced from the wintering area. *Ethology Ecology, Evolution*, 7: 257-264.
- BACCETTI N., DALL'ANTONIA P., MAGAGNOLI P., MELEGA L., SERRA L., SOLDATINI C., ZENATELLO M., 2002. Risultati dei censimenti degli uccelli acquatici svernanti in Italia: distribuzione, stima e trend delle popolazioni nel 1991 – 2000. *Biol. Cons. Fauna* 111:1-240.
- BACCETTI N., SERRA L., CHERUBINI G., 1999. Timing of attachment to wintering site as revealed by experimental displacement of Dunlins (*Calidris alpina*). *J. Ornithol.*, 140: 309-317.
- BACCETTI N., SERRA L., TINARELLI R., UTMAR P., CHERUBINI G., KRAVOS K., CASINI L., 1992. Nuovi conteggi di limicoli costieri svernanti nelle zone umide adriatiche. *Riv. ital. Orn.*, 62:3-12.
- BAGLIACCA M. Gestione delle specie aviarie in difficoltà: alimentazione e relativo comportamento gestionale–L.I.P.U.–C.R.U.M.A–Livorno-in: <http://www.vet.unipi.it/new/didattica/stpa/ecofauna0405/Bagliacca/lipu.pdf>
- BAIRLEIN F., 2003. The study of bird migrations – some future perspectives. *Bird Study* 50: 243 – 253.
- BAKER A.J., 2002. The deep roots of bird migration: inferences from the historical record preserved in DNA. *Ardea* 90: 503 – 513.

- BAKER A.J., PIERSMA T., ROSENMEIER L., 1994. Unraveling the intraspecific phylogeography of knots *Calidris canutus*: a progress report on the search for genetic markers. *J Ornithol* 135: 599 – 608.
- BARRET R. T., CAMPHUYSEN K. (C.J.), ANKER-NILSSEN T., CHARDINE J. W., FURNESS R. W., GARTHE S., HUPPOP O., LEOPOLD M.F., MONTEVECCHI W.A., VEIT R.R., 2007. *Journal of Marine Science Advance Access*, 1-17.
- BAUCHINGER U. & BIEBACH H., 2001. Differential catabolism of muscle protein in Garden Warblers (*Sylvia borin*): flight and leg muscle act as a protein source during long-distance migration. *J Comp Physiol B*, 171: 293-301.
- BELL D.V. & OWEN M., 1990. Shooting disturbance-A review. In: Matthews, G.V.T. (Ed); Managing waterfowl populations. *IWRB Spec. Pub.* 12. IWRB, Slimbridge: 159-171.
- BERETTA C., 1994. Tossicologia Veterinaria. Casa Editrice Ambrosiana, Milano.
- BERTHOLD P., 2003. La migrazione degli uccelli. Bollati Boringhieri editore s.r.l., Torino.
- BEYER H.L., 2004. Hawth's Analysis Tools for ArcGIS. <http://www.spatial ecology.com/htools>.
- BIANCHI C.N., ARDIZZONE G.D., BELLUSCIO A., COLANTONI P., DIVIACCO G., MORRI C., TUNESI L., 2003. La cartografia del benthos. *Biol. Mar. Medit.*, 19 (suppl.): 367-394
- BIBBY C.J., BURGESS N.D., HILL D.A., MUSTOE S.H., 2000. Bird census techniques. 2nd edition. Academic Press, 302+XVI pp.
- BINO T. & JORGO G., 2002. Conservation status and threats over Albanian wetlands. Albanian Society for the Protection of birds and Mammals. Relazione inedita, 47 pp.
- BINO T., TOURENQ C., KAYSER Y., BUSUTTI S., CROZIER J., DORE B.J., BEGO F., 1996. Recensement des oiseaux d'eau hivernants en Albanie (14 -31 janvier 1996). Station Biologique de la Tour du Valat and Natural History Museum, Tirana. Relazione inedita, 102 pp.
- BLANC R., GUILLEMAIN, M., MOUNRONVAL, J.B., DESMONTS, D., FRITZ, H., 2006. Effects of nonconsumptive leisure disturbance to wildlife. *Revue d'Ecologie (Terre, Vie)* 61: 117-133.
- BÖHNING-GÄSE K., SCHUDA M.D., HELBIG A.J., 2003. Weak phylogenetic effects on ecological niches of *Sylvia* warblers. *J Evol Biol* 16: 956 – 965.
- BOOM R., SOL C.J., SALIMANS M.M., JANSEN C.L., WERTHEIM-VAN DILLEN P.M., VAN DER N.J., 1990. Rapid and simple method for purification of nucleic acids. *Journal of Clinical Microbiology* 28: 495-503.
- BOTH K., SLY N., BURGOYNE L., 2000. FTA paper, DNA, time and the profiler. In: <http://129.96.230.55/vidocq>.
- BOTTO F. & IRIBARNE O., 2000. Contrasting effects of two borrowing carbs (*Chasmagnathus granulata* and *Uca uruguayensis*) on sediment composition and transport in estuarine environments. *Estuarine, Coastal and Shelf Science*, 51: 141-151.
- BOUDOURESQUE C.F., 1971. Méthodes d'études qualitative et quantitative du benthos. *Tethys*, 3: 79-104.
- BRAMWELL R.K., 2003. Sexing chicks in the backyard flock. *Avian Advice* 5: 4 – 5.
- BRICHETTI P. & FRACASSO G., 2003. Ornitologia italiana. Vol.1- Gaviidae-Falconidae. Alberto Perdisa editore, Bologna.
- BRICHETTI P. & FRACASSO G., 2004. Ornitologia Italiana. Vol. 2 – Tetraonidae-Scolopacidae. Alberto Perdisa Editore, Bologna.
- BUCHLOLZ W.G., PEARCE J.M., PIERSON B.J., SCRIBNER K.T., 1998. Dinucleotide repeat polymorphisms in waterfowl (family Anatidae): characterization of a sex-linked (Z-specific) and 14 autosomal loci. *Anim. Genet.* 29: 323 – 325.

- BUIA M.C. & MAZZELLA L., 1991. Reproductive strategies of the Mediterranean seagrasses: *Posidonia oceanica* (L.) Delile, *Cymodocea nodosa* (Ucria) Ascher and *Zostera noltii* (Hornem). *Aquat. Bot.*, 40: 343-362.
- BUIA M.C., RUSSO G.F., MAZZELLA L., 1985. Interrelazioni tra *Cymodocea nodosa* e *Zostera noltii* in un prato misto superficiale dell'isola di Ischia. *Nova Thalassia*, 7: 406-408.
- BURGER J., 1993. Metals in avian feathers: bioindicators of environmental pollution. *Reviews in Environmental Toxicology* 5, pp. 203-311.
- BURGOYNE L. & ROGERS C., 1997. Bacterial typing: storing and processing of stabilized reference bacteria for polymerase chain reaction without preparing DNA: an example of an automatable. *Procedure of Analytical Biochemistry*, 247, 223 – 227.
- CADENNE, 1981. Citato in: <http://veterinaria.campusnet.unito.it/didattica/att/a68c.1997.file.pdf>
- CAMPREDON P., 1981. Hivernage du canard siffleur *Anas penelope* L. en Camargue (France). Stationnements et activités. *Alauda* 49: 161-193.
- CAMPREDON S., CAMPREDON P., PIROT J.Y., TAMISIER A., 1982. Manuel d'analyse des contenus stomacaux de canard et de foulques. Centre d'Ecologie de Camargue. Office National de la Chasse. 87 pp.
- CANIGLIA G., BORELLA S., CURIEL D., NASCIMBENI P., PALOSCHI F., RIMONDO A., SCARTON F., TAGLIAPIETRA D., ZANELLA L., 1990. Cartografia della distribuzione delle fanerogame marine nella laguna di Venezia. *Giorn. Bot. Ital.*, 124(1): 212.
- CARESSA S., CESCHIA C., OREL G., TRELEANI R., 1995. Popolamenti attuali e pregressi nel Golfo di Trieste da Punta Salvore a Punta Tagliamento (Alto Adriatico). In: Cinelli F., Fresi E., Lorenzi C., Mucedola A. *La Posidonia oceanica. Un contributo per la salvaguardia del principale ecosistema marino del Mediterraneo. Rivista marittima*: 160-187.
- CASINI L., MAGNANI A., SERRA L., 1992. Ciclo annuale della comunità degli uccelli acquatici nella Salina di Cervia. *Ric. Biol. Selvaggina* 92: 1-54.
- CAYE G. & MEINESZ A., 1985. Observation on the vegetative development, flowering and seedling of *Cymodocea nodosa* (Ucria) Ascher. On the Mediterranean coasts of France. *Aquat. Bot.*, 22: 277-289.
- CAYFORD, J., 1993. Wader disturbance: A theoretical overview. *Wader Study Group Bulletin* 68: 3-5.
- CIRAOLO G., CALVO S., LA LOGGIA G., 2003. Monitoring *Posidonia oceanica* meadows in a mediterranean coastal lagoon (Stagnone, Italy) by means of neural network and isodata classification methods. *Int. J. Remote Sens.*, 24(13): 2703-2716.
- CISO-COI, 2007. Lista CISO-COI. Check-list degli Uccelli (Aves) italiani aggiornata al 25-01-2005. Pagina web: <http://www.ciso-coi.org/COImateriale/ListaCISO-COI.pdf> (Query date: 17/09/2008)
- CLENCH M. H. & MATHIAS R. J., 1995. The avian cecum: a review. *Wilson Bull.*, 107(1), pp 93-121
- COHEN A. N., CARLTON J. T., FOUNTAIN M. C., 1995. Introduction, dispersal and potential impacts of the green crab *Carcinus maenas* in San Francisco bay, California. *Marine Biology*, 122: 225-237.
- COHEN J., 1960. A coefficient of agreement for nominal scales. *Educ. Psychol. Meas.* 20:37-46
- COSOLO M., UTMAR P., ROPPA F., SPONZA S., 2007. Ruolo ecologico delle zone umide costiere del Friuli Venezia Giulia per l'avifauna acquatica: importanza delle zone di marea e delle valli arginate. Linee guida e casi di studio per la gestione dei siti della rete Natura 2000 in ambienti di transizione; atti del Workshop di Grado 7-8 giugno 2006. Univ. di Trieste: 197 – 212.

- COSSIGNANI T., COSSIGNANI P., DI NISIO A., PASSAMONTI M., 1992. Atlante delle conchiglie del Medio Adriatico. Mostra Mondiale Malacologia Cupra Marittima (AP, Italy). L'Informatore Piceno. Ed. Ancona: 223.
- COTTIGLIA M., 1983. Decapodi lagunari. Monotipia erredi – Genova, pp. 50 – 52, 135 – 137.
- CRAMP S. & SIMMONS K.L.S., 1977. The birds of the Western Palearctic. 1. Ostrich to Ducks. Oxford University Press, Oxford.
- CRONIN M.A., GRAND J.B., ESLER D., DERKSEN D.V., SCRIBNER K.T., 1996. Breeding populations of Northern pintails have similar mitochondrial DNA. *Can J Zool* 74: 992 – 999.
- D'ANGELO G. & GARGIULLO S., 1979. Guida alle conchiglie mediterranee. Conoscerle, cercarle, collezionarle. Fabbri Editore.
- DAUWE T., BERVOEST L., JANSSENE E., PINXTEN R., BLUST R., EENS M., 2002. Great and blue tit feathers as biomonitors for heavy metal pollution. *Ecological indicators* 1, pp.227-234.
- DAVIDSON N.C. & ROTHWELL P.I., 1993b. Disturbance to waterfowl on estuaries: The conservation and coastal management implications of current knowledge. *Wader Study Group Bulletin* 68: 97-105.
- DAWSON T. J., WHITEHEAD P. J., MCLEAN A., FANNING F. D AND R. DAWSON W., 2000. Digestive function in Australian magpie geese (*Anseranas semipalmata*). *Australian Journal of Zoology* 48(3) 265 – 279.
- DAY J.W., 1992. Sea level rise, management options and the future of Mediterranean coastal wetlands. in *Managing Mediterranean wetlands and their birds*, IWRB Special Publication No. 20.
- DEL RIO S., 2000. Cost-effectiveness in sample processing using the FTA[®] treated stain card for high throughput. In: <http://www.promega.com>.
- DELANY, S. & SCOTT, D.A. 2006. - Waterbird Population Estimates; Fourth Edition. Wetlands International, Wageningen, The Netherlands
- DEMAREST D.W., KAMINSKY R.M., BRENNAN L.A., BOYLE C.R., 1997. Body-mass, survival, and pairing consequences of winter-diet restriction wood ducks. *J. Wildl. Manage.* 61:822-832.
- DEMETRESCU C., FINOCCHI I., ITALIANO G.F., 2004. Algoritmi e strutture dati McGraw-Hill, Milano, ISBN 88-386-6161-8, pp. 447
- DENHARTOG C., 1970. The seagrass of the world. Amsterdam, Verh. Kon. Ned. Akad. Wet., Afd. *Natuurk*: 275.
- DERMANIS A. & BIAGI L., 2002. Telerilevamento. Informazione territoriale mediante immagini da satellite. Casa Editrice Ambrosiana. Milano, pp. 179-194.
- DIETZ M.V., DE KINGA A., PIERSMA T. AND VERHULSTS S., 1999. Estimating organ size in small migrating shorebirds with ultra-sonography: an inter-calibration exercise. *Physiological and biochemical zoology* 72, pp.28-37.
- DONNE-GOUSSE C., LAUDET V., HÄNNI C., 2002. A molecular phylogeny of anseriformes based on mitochondrial DNA analysis. *Mol. Phylogenet. Evol.* 23: 339 – 356.
- DUARTE C.M., KIRKMAN H., 2001. Methods for measurement of seagrass abundance and depth distribution. In: Short F.T., Coles R.G.(eds). *Global seagrass research methods*. Elsevier, Amsterdam:141-153.
- DUBOVSKY J.A. & KAMINSKY R.M., 1994. Potential reproductive consequence of winter-diet restriction in mallards. *J. Wildl. Manage.* 58: 780-786.
- DURAND, F., DEVILLERS, N., LALLIER, F.H., REGNAULT, M., 2000. Nitrogen excretion and changes in blood components during emersion of the subtidal spider crab *Maia squinado* (L). *Comparative Biochemistry and Physiology*, 127A: 259–271.

- DURANT D., 2003. The digestion of fibre in herbivorous Anatidae: a review. *Wildfowl* 54 7–24.
- DURANT D., FRITZ H., BLAIS S., 2003. The functional response in three species of herbivorous Anatidae: effects of sward height, body mass and bill size. *Journal of Animal Ecology* n° 72 pp 220-231.
- EASTMAN J.R., 2001. IDRISI for Windows, version IDRISI32 Release 2, Worcester MA: Clark school of Geography, Clark University.
- EGGLESTONE D.B., ETHERINGTON L.L., WARD E.E., 1998. Organism response to habitat patchiness: species and habitat- dependent recruitment of decapod crustaceans. *Journal of Experimental Marine Biology and Ecology*, 223: 111–132.
- EISLER R., 1986. Chromium hazard to fish, wildlife and invertebrates: a synoptic review. Patuxent Wildlife Research Center, Biological Report n°6.
- EISLER R., 1988a. Arsenic hazard to fish, wildlife and invertebrates: a synoptic review. Patuxent Wildlife Research Center, Biological Report n°12.
- EISLER R., 1988b. Lead hazard to fish, wildlife and invertebrates: a synoptic review. Patuxent Wildlife Research Center, Biological Report n°14.
- ELLEGREN H., 1996. First gene on the avian W chromosome (CHD) provides a tag for universal sexing in non-ratide birds. *Proc. Royal Soc. London B* 263: 1635 – 1641.
- ELLEGREN H., 2001. Hens, cocks and avian sex identification. A quest for genes on Z or W. *EMBO Reports* 2: 192 – 196.
- EUROPEAN ENVIRONMENTAL AGENCY, 2003. Europe's environment: the third assessment. Environmental assessment report No. 10.
- EVANS D.M. & DAY K.R., 2001. Does shooting disturbance affect diving ducks wintering on large shallow lakes? A case study on Lough Neagh, Northern Ireland. *Biological Conservation* 98: 315-323.
- FACCHIN G. & FLORIT F., 2005. Progetto di censimento degli uccelli acquatici svernanti in Friuli Venezia Giulia nell'ambito dell'*International Waterfowl Census* (IWC). Relazione sintetica ad uso interno sull'attività dell'anno 2005. Regione Autonoma Friuli Venezia Giulia - Direzione centrale risorse agricole, naturali, forestali e montagna - Servizio tutela ambienti naturali, fauna e Corpo forestale regionale - Ufficio studi faunistici
- FACCHIN G., FLORIT F., ZENATELLO M., *in stampa*. La banca dati delle zone umide soggette al censimento degli uccelli acquatici svernanti (*International Waterbird Census* – IWC) in Friuli Venezia Giulia. XIV Convegno Italiano di Ornitologia. Trieste, 26-30 settembre 2007
- FARINHA J.C., COSTA L.T., ZALIDIS G., MANTZAVELAS A., FITOKA E., HECKER N., TOMÁS VIVES P., 1996. Mediterranean Wetland Inventory: Data Recording. MedWet. Instituto da Conservação da Natureza. Wetlands International publication. Volume III.
- FIELDS R.L. & SCRIBNER K.T., 1997. Isolation and characterization of novel waterfowl microsatellite loci: cross-species comparisons and research applications. *Mol. Ecol.* 6: 199 – 202.
- FOCKEDEVY N., MEES J., VANGHELUWE M., VERSLIJCKE T., JANSSEN C.R., VINCX M., 2005. Temperature and salinity effects on post-marsupial growth of *Neomysis integer* (Crustacea: Mysidacea). *Journal of Experimental Marine Biology and Ecology*, 326: 27-47.
- FOX A.D. & MADSEN J., 1997. Behavioural and distributional effects of hunting disturbance on waterbirds in Europe: implication for refuge design. *Journal of Applied Ecology* 34: 1-13.
- FRIDOLFSSON A-K. & ELLEGREN H., 1999. A simple and universal method for molecular sexing of non-ratide birds. *J. Avian Biol.* 20: 116 – 121.
- GHIRARDELLI E., 1981. La vita nelle acque. Il nostro universo, UTET.

- GILBERT G., GIBBONS D.W., EVANS J., 1998. Bird Monitoring Methods. RSPB, 464 pp.
- GILL J.A., 2007. Approaches to measuring the effects of human disturbance on birds. *Ibis* 149(1): 9-14.
- GILL J.A., NORRIS, K., SUTHERLAND, W.J., 2001a. Why behavioural responses may not reflect the population consequences of human disturbance. *Biological Conservation* 97: 265-268.
- GILL J.A., SUTHERLAND, W.J., WATKINSONS, A.R., 1996. A method to quantify the effects of human disturbance on animal populations. *Journal of Applied Ecology* 33: 786-792.
- GILLIKIN D.P, DE WACHTER B., TACK J.F., 2004. Physiological responses of two ecologically important Kenyan mangrove crabs exposed to altered salinity regimes. *Journal of Experimental Marine Biology and Ecology*, 301: 93-109.
- GLENN T.C., THOMPSON J.E., BALLARD B.M., ROBERSON J.A., FRENCH J.O., 2002. Mitochondrial DNA variation among wintering midcontinent gulf coast sandhill cranes. *J Wildl Manage* 66: 339 – 348.
- GOERING, H.K. & VAN SOEST, P.J., 1970. Forage fibre analyses (apparatus, reagent, procedures and some applications). US Department of Agriculture, Agriculture Handbook, 379 pp.
- GORDON R.F., 1981. Patologia Aviare. Ermes, Milano.
- GOSS-CUSTARD J.D. & DURRELL S.E.A., 1990. Bird behaviour and environmental planning: approaches in the study of wader populations. *Ibis*, 132: 273-289.
- GOSS-CUSTARD J.D., CALLOW R.W.G., CLARKE R.T., 1992. Correlates of the density of foraging oystercatchers *Haematopus ostralegus* at different population sizes. *J. Anim. Ecol.*, 61: 159-173.
- GOSS-CUSTARD, J.D., STILLMAN, R.A., WEST, A.D., CALDOW, R.W.G., MCGRORTY, S., 2002. Carrying capacity in overwintering migratory birds. *Biological conservation* 105: 27-41.
- GREEN E.P. & SHORT F.T., 2003. World Atlas of Seagrass. Univ. California press.(USA).
- GRIFFITHS R. & KORN R., 1997. A CHD1 gene is Z chromosome linked in the Chicken *Gallus domesticus*. *Gene* 197: 225 – 229.
- GRIFFITHS R. & TIWARI B., 1993. The isolation of molecular genetic markers for the identification of sex. *Proceedings of the National Academy of Sciences* 90: 8324 – 8326.
- GRIFFITHS R. & TIWARI B., 1995. Sex of the last wild Spix's macaw. *Nature* 375: 454.
- GRIFFITHS R., DOUBLE M., ORR K., DAWSON R., 1998. A DNA test to sex most birds. *Mol. Ecol.* 7: 1071 – 1075.
- GUELORGET O. & PERTHUISOT J.P., 1983. Le domaine paralique. Expressions géologiques, biologiques et économiques du confinement. Travaux du laboratoire de géologie, 16.
- GUERZONI S. & TAGLIAPIETRA D. (a cura di), 2006. Atlante della laguna di Venezia. Ed. Marsilio.
- GUGLIELMO C.G., CERASALE D.J., ELDERMIRE C., 2005. A field validation of plasma metabolite to assess refuelling performance of migratory birds. *Physiological and Biochemical Zoology* 78 (1), pp.116-125.
- GUGLIELMO C.G., O'HARA P.D., WILLIAMS T.D., 2002. Extrinsic and intrinsic sources of variation in plasma lipid metabolites of free-living Western Sandpipers (*Calidris mauri*). *The Auk* 119, pp. 437-445.
- GUILLEMAIN M., FRITZ H., DUNCAN P., 2002. The importance of protected areas as nocturnal feeding grounds for dabbling ducks wintering in western France. *Biological Conservation* 103: 183-198.
- GUSTIN M., ROSSI P., CELADA C., 2005. Aggiornamento delle conoscenze ornitologiche nelle IBA (*Important Bird Areas*) e sviluppo di proposte tecnico – scientifiche sul completamento della rete di ZPS (Zone di Protezione Speciale) in Friuli Venezia Giulia. Relazione ad uso della Regione autonoma Friuli Venezia Giulia. Pp. 71.

- GUZZON C. & PANZARIN L., 2005. Forapaglie comune *Acrocephalus shoenoaenus* nidificante in cariceti soggetti a marea dell'Alto Adriatico: siti riproduttivi nuovi o ritrovati dopo un secolo? *Avocetta* 29: 81.
- GUZZON C. & UTMAR P., 2005. The importance of protected areas for breeding and wintering waterbirds in the coastal wetlands of Friuli-Venezia Giulia (North-East Italy). In: ASOER (ed.) - Avifauna acquatica: esperienze a confronto. Atti del I Convegno (30 aprile 2004, Comacchio). Tipografia Giari, Codigoro, pp. 66-67.
- GUZZON C., KRAVOS K., UTMAR P., 2001. Censimenti mensili di limicoli nelle zone umide costiere del Friuli-Venezia Giulia. Primi dati: anno 1999. In TELLINI FLORENZANO G., BARBAGLI F., BACCETTI N., (Eds.). Atti XI Conv. It. Orn., *Avocetta*, 25: 217.
- GUZZON C., TOUT C.P., UTMAR P. (a cura di), 2005. I censimenti degli uccelli acquatici svernanti nelle zone umide del Friuli Venezia Giulia, anni 1997-2004. Associazione Studi Ornitologici e Ricerche Ecologiche del Friuli Venezia Giulia (A.ST.O.R.E.-FVG). 'Centro Stampa' di A. Candido, F. Spanghero Snc – Monfalcone (GO).
- HAGEMEIJER J.M. & BLAIR M.J. (eds.), 1997. The EBCC Atlas of European Breeding Birds. Their Distribution and Abundance. T., A.D. Poyser, London, 903 pp.
- HAIG S.M., MULLINS T.D., FORSMAN E.D., TRAIL P.W., WENNERBERG L., 2004. Genetic identification of Spotted Owls, Barred Owls and their hybrids: legal implications of hybrid identity. *Conserv Biol* 18: 1347 – 1357.
- HALL D.J., & BALL G.B., 1965. *ISODATA: a novel method of data analysis and pattern classification*. Technical report, Stanford Research Institute, Menlo Park, CA.
- HESTBECK J.B., 1993. Overwinter distribution of northern pintail population in North America. *J. Wildl. Manage.* 57: 582-589.
- HOLLAMBY S., AFEMA-AZIKURU J., WAIGO S., CAMERON K., RAE GANDOLF A., NORRIS A. AND SIKARSKIE J. G., 2006. Suggested Guidelines for Use of Avian Species as Biomonitors. *Environmental Monitoring and Assessment* 118 (1-3), pp. 13-20.
- HOLLIS G.E., 1986. The Modelling and Management of the Internationally Important Wetland at Garaet El Ichkeul, Tunisia , IWRB Special Publication No 4.
- HUANG G.H. & CHANG N.B., 2003. Perspectives of Environmental Informatics and Systems Analysis *Journal of Environmental Informatics* Vol 1, pp. 1-6
- HUANG Y., TU J., CHENG X., TANG B., HU X., LIU Z., FENG J., LOU Y., LIN L., XU K., ZHAO Y., LI N., 2005. Characterization of 35 novel microsatellite DNA markers from the duck (*Anas platyrhynchos*) genome and cross-amplification in other birds. *Genet. Sel. Evol.* 37: 455 – 472.
- HUDSON W.D. & RAMM C.W., 1987. Correct Formulation of the Kappa Coefficient of Agreement. *Photogramm. Eng. Remote Sens.*, 53(4): 421-422.
- INFS, 1999. Regolamento per lo svolgimento dell'attività di inanelamento a scopo scientifico. Litografia F.G. Savignano s. P. – Modena
- IRIBARNE O., BRUSCHETTI M., ESCAPA M., BAVA J., BOTTO F., GUTIERREZ J., PALOMO G., DELHEY K., PETRACCI P., GAGLIARDINI A., 2005. Small- and large- scale effect of the SW Atlantic Burrowing crab *Chasmagnathus granulatus* on habitat use by migratory shorebirds. *Journal of Experimental Marine Biology and Ecology*, 315: 87-101.
- IRIBARNE O., MARTINEZ M.M., 1999. Predation of the southwestern atlantic fiddler crab (*Uca uruguayensis*) by migratory birds (*Pluvialis dominica*, *P. squatarola*, *Arenaria interpres*, and *Numenius phaeopus*). *Estuaries*, 22: 47-54.

- JAX K., 2006. Ecological Units: definitions and application. *The Quarterly Review of Biology*, 81, 3, 237-258.
- JENSEN J.R., 1986. *Introductory Digital Image Processing*. Prentice-Hall, Englewood Cliffs, New Jersey, pp. 379.
- JOHNSON K.P. & SORENSON M.D., 1998. Comparing molecular evolution in two mitochondrial protein coding genes (cytochrome b and ND2) in the dabbling ducks (Tribe: Anatini). *Mol. Phylogenet. Evol.* 10: 82 – 94.
- JOHNSON K.P. & SORENSON M.D., 1999. Phylogeny and biogeography of dabbling ducks (Genus: *Anas*): a comparison of molecular and morphological evidence. *The Auk* 116: 792 – 805.
- JORDE D.G., KRAPU G.L., CRAWFORD R.D., 1983. Feeding ecology of Mallards wintering in Nebraska. *Journal of Wildlife Management* 47: 104-1053.
- JOSEPH L., WILKE T., ALPERS D., 2003. Independent evolution of migration on the South American landscape in a long-distance temperate-tropical migratory bird, Swainson's flycatcher (*Myiarchus swainsoni*). *J Biogeogr* 30: 925 – 937.
- JÓZEFIAK D., RUTKOWSKI A., MARTIN S. A., 2004. Carbohydrate fermentation in the avian ceca: a review - *Animal feed science and technology*. Vol.113, n.1-4, pp. 1-15. Elsevier.
- KAHN N.W., JOHN J., QUINN T., 1998. Chromosome-specific intron size differences in the Avian CHD gene provide an efficient method for sex identification in birds. *The Auk* 115: 1074 – 1078.
- KAYSER Y., BINO T., GAUTIER-CLERC M., 1995. Recensement des oiseaux d'eau hivernants en Albanie 17 janvier – 7 février 1995. Station Biologique de la Tour du Valat. Relazione inedita, 79 pp.
- KAYSER Y., BINO T., BEGO F., FREMUTH W., JORGO G., 1997. Recensement des oiseaux d'eau hivernants en Albanie (3- 19 janvier 1997). Station Biologique de la Tour du Valat and Natural History Museum, Tirana. Relazione inedita, 52 pp.
- KENWARD R.E., 2001. *A manual for wildlife radio tagging*. Academic Press, London
- KIMURA M., CLEGG S.M., LOVETTE I.J., HOLDER K.R., GIRMAN D.J., MILA B., WADE P., SMITH T.B., 2002. Phylogeographical approaches to assessing demographic connectivity between breeding and overwintering regions in a nearctic-neotropical warbler (*Wilsonia pusilla*). *Mol* 11: 1605 – 1616
- KLASING K. C., 1998. *Comparative Avian Nutrition 1998*. CAB International.
- KRAVOS K., CANDOTTO S., GUZZON C., UTMAR P., 2003. La presenza del Fischione *Anas penelope* nella fascia costiera del Friuli-Venezia Giulia (1989-2002). In: Conti P., Ruolini D., Galeotti P., Milone M., De Filippo G. (a cura di). *Atti XII Conv. It. Orn., Avocetta* 27: 163.
- KULIKOVA I.V., ZHURAVLEV Y.N., MCCracken K.G., 2004. Asymmetric hybridization and sex-biased gene flow between Eastern Spotbilled Ducks (*Anas zonorhyncha*) and Mallards (*A. platyrhynchos*) in the Russian Far East. *The Auk* 121: 930 – 949.
- KUO J., DENHARTOG C., 2001. Seagrass taxonomy and identification key. In: Short F., Coles R. (eds). *Global seagrass research methods*. Elsevier, Amsterdam: 31-58.
- KVIST L., RUOKONEN M., LUMME J., ORELL M., 1999. The colonization history and present-day population structure of the European great tit (*Parus major major*). *Heredity* 82: 495 – 502.
- LACERDA T.P. & SAWAYA P., 1986. Effects of hypo – osmotic conditions on haemolymph glucose concentration in *Callinectes danae* Smith 1869 (Crustacea, Decapoda, Portunidae). *Comparative Biochemistry and Physiology*, 85A: 509–512.
- LANDIS J.R. & KOCH G.G., 1977. The measurements of observer agreement for categorical data. *Biometrics* 33: 159-174

- LANDYS M.M., PIERSMA T., GUGLIELMO C.G., JUKEMA J., RAMENOFKY M., WINGFIELD J.C., 2005. Metabolic profile of long-distance migratory flight and stopover in a shorebird. *Proceedings of the Royal Society B*.
- LEGENDRE L. & LEGENDRE P., 1983. Numeral ecology. Developments in environmental modelling. *Elsevier Scient. Publ. Co., Amsterdam*. XVI. 419
- LEICA GEOSYSTEMS ERDAS IMAGINE 8.7. GIS AND MAPPING, 2003. Atlanta, Georgia, USA, Leica Inc.
- LIEBERS D., HELBIG A.J., DE KNIJFF P., 2001. Genetic differentiation and phylogeography of gulls in the *Larus cachinnans-fuscus* group (Aves: Charadriiformes). *Mol Ecol* 10: 2447 – 2462.
- LILLEBO A., PARDAL M.A., NETO J. M., MARQUES J.C., 2003. Salinity as the major factor affecting *Scirpus maritimus* annual dynamics. Evidence from field data and greenhouse experiment. *Aquatic Botany*, 77: 111-120.
- LINDSTROM A., KVIST A., PIERMA T., DEKINGA M.W., 2000. Avian pectoral muscle size rapidly tracks body mass changes during flight, fasting and fuelling. *The Journal of Experimental Biology* 203, pp. 913-919.
- LORENZON S., GIULIANINI P.G., LIBRALATO S., MARTINIS M., FERRERO E.A., 2008. Stress of two different transport systems on the physiological profile of the crab *Cancer pagurus*. *Aquaculture*, 278: 156-163.
- LORENZON S., GIULIANINI P.G., MARTINIS M., FERRERO E.A., 2007. Stress effect of different temperatures and air exposure during transport on physiological profiles in the American lobster *Homarus americanus*. *Comparative Biochemistry and Physiology, Part A*, 147: 94-102.
- LOVETTE I.J., CLEGG S.M., SMITH T.B., 2004. Limited utility of mtDNA markers for determining connectivity among breeding and overwintering locations in three neotropical migrant birds. *Conserv Biol* 18: 156 – 166.
- LUIS A. & GOSS-CUSTARD J.D., 2005. Spatial organization of the Dunlin *Calidris alpina* L. during winter – the existence of functional units. *Bird Study*, 52: 97-103.
- MAAK S., WIMMERS K., WEIGEND S., NEUMANN K., 2003. Isolation and characterization of 18 microsatellites in the Peking duck (*Anas platyrhynchos*) and their application in other waterfowl species. *Mol. Ecol. Notes* 3: 224 – 227.
- MADSEN J., 1995. Impacts of disturbance on migratory waterfowl. *Ibis* 137: S67-S74.
- MADSEN J., 1998a. Experimental refuges for migratory waterfowl in Danish wetlands. I. Baseline assessment of the disturbance effects of recreational activities. *Journal of Applied Ecology* 35: 386-397.
- MADSEN J., 1998b. Experimental refuges for migratory waterfowl in Danish wetlands. II. Tests of hunting disturbance effects. *Journal of Applied Ecology* 35: 398-417.
- Manuale dell'Operatore SonoAce PICO, Medison.
- MARBA N., DUARTE C.M., 1997. Interannual changes in seagrass (*Posidonia oceanica*) growth and environmental change in the Spanish Mediterranean littoral zone. *Limnol. Oceanogr.*, 42: 800-810.
- MARTINEZ-CRUZ B., GODOY J.A., NEGRO J.J., 2004. Population genetics after fragmentation: the case of the endangered Spanish imperial eagle (*Aquila adalberti*). *Mol Ecol* 13: 2243 – 2255.
- MATHERS R.G. & MONTGOMERY W.I., 1997. Quality of food consumed by over wintering pale bellied Brent Geese *Branta Bernicla Hrota* and Wigeon *Anas Penelope*. *Biology and Environment: Proceedings of the Royal Irish Academy*. Vol. 97B, n° 2, 81-89.

- MATHEVET R. & TAMISIER A., 2002. Creation of a nature reserve, its effect on hunting management and waterfowl distribution in the Camargue (southern France). *Biodiversity and Conservation* 11: 509-519.
- MAYHEW P.W., 1988. The daily energy intake of European Wigeon in winter. *Ornis Scand.* 19: 217-223.
- MCKELVEY R.W. & SMITH G.E.J., 1990. The distribution of waterfowl banded or returned in British Columbia, 1951-1985. *Can Wildl. Serv. Tech. Rep. Ser.*, 79. Delta, British Columbia.
- MILA B., GIRMAN D.J., KIMURA M., SMITH T.B., 2000. Genetic evidence for the effect of a postglacial population expansion on the phylogeography of a North American songbird. *Proc R Soc Lond – B* 267: 1033 – 1040.
- MILOT E., GIBBS H.L., HOBSON K.A., 2000. Phylogeography and genetic structure of Northern populations of the yellow warbler (*Dendroica petechia*). *Mol Ecol* 9: 667 – 681
- MIYAKI C.Y., DUARTE M.B., CAPARROZ R., NUNES A.V., WAJNTALA., 1997. Sex identification of South American Parrots (Psittacidae, Aves) using the human minisatellite probe 33.15. *The Auk* 114: 516 – 520.
- MOLINIER R. & PICARD J., 1952. Reserches sur les herbiers de phanerogames marines du littoral méditerranéen français. *Ann. Inst. Oceanogr.*, 27(3): 127-134.
- MOREIRA F., 1997. The importance of shorebirds to energy fluxes in a food web of a south European estuary. *Estuarine, Coastal and Shelf Science*, 44: 67-78.
- MOSS R., HANSSSEN I., 1980. Grouse nutrition. *Nutrition abstract and reviews*, 50 (11). 555-567.
- MÚRIAS, T., CABRAL J.A., LOPES R., MARQUES J.C., GOSS-CUSTARD J.D., 2002. Use of traditional salines by waders in the Mondego estuary (Portugal): a conservation perspective. *Ardeola*, 49: 223-240.
- MYERS J.P., 1984. Spacing behaviour of nonbreeding shorebirds, p.27 1-32 1. In: Burger J. and B.L. Olla [eds.]. *Shorebirds: Migration and Foraging Behaviour*. Plenum Press, New York.
- NACKEN M. & KARSTEN R., 2000. Effects of herbivorous birds on intertidal seagrass beds in the northern Wadden Sea. *Helgoland Marine Research*, 54: 87–94.
- NAS, 1978. An assessment of mercury in the environment. United States National Academy of Sciences, Washington, DC, 185
- NEGRONI G., 2000. Final report T.A. to F.D. PHARE project, 2000.
- NESJE M. & RØED K., 2000. Sex identification in falcons using microsatellite DNA markers. *Hereditas* 132: 261 – 263.
- NEWTON S.F., 1993. Body condition of a small passerine bird: Ultrasonic assessment and significance in over-winter survival. *J. Zool. Lond.* 229: 561–580.
- NOVO, M. S., MIRANDA R. B., BIANCHINI A., 2005. Sexual and seasonal variation in osmoregulation and ionoregulation in the estuarine crab *Chasmagnathus granulatus* (Crustacea, Decapoda). *Journal of Experimental Marine Biology and Ecology*, 323: 118-137.
- OCAMPO, L., PATINO, D., RAMÍREZ, C., 2003. Effect of temperature on hemolymph lactate and glucose concentrations in spiny lobster *Panulirus interruptus* during progressive hypoxia. *Journal of Experimental Marine Biology and Ecology*, 296: 71–77.
- ODORICO R. & BRESSAN G., 1992. Variazioni nella fisionomia di aree a fanerogame marine (Riserva Marina di Miramare-Trieste). *Boll. Soc. Adriatica Scienze*, 73: 1-61.
- OREL G., 1988. Aspetti della bionomia bentonica e della pesca nel Golfo di Trieste con particolare riferimento ai fondali prospicienti il promontorio di Miramare. *Hydrores*, anno V, 6: 57-70.
- OWEN M., 1973. The winter feeding ecology of Wigeon at Bridgwater Bay, Somerset. *Ibis* 115: 227-243.

- OWEN M. & THOMAS G., 1979. The feeding ecology and conservation of Wigeon wintering at the Ouse Washes, England. *J. Appl. Ecol.* 16: 765-809.
- PAILLISSON J.M., REEBER S., MARION L., 2002. Bird assemblage as bio-indicators of water regime management and hunting disturbance in natural wet grasslands. *Biological conservation*, 106(1): 115-127.
- PALOMO G., BOTTO F., NAVARRO D., ESCARA M., IRIDARNE O., 2003. Does the presence of the Sw Atlantic borrowing crab *Chasmagnathus granulatus* Dana affect predator – prey interactions between shorebirds and polychaetes? *Journal of Experimental Marine Biology and Ecology*, 290: 211-228
- PARENZAN P., 1974. Carta d'identità delle conchiglie del Mediterraneo. Ed. Bios Taras. Vol. I.e II.
- PARODI R. & PERCO F., 1988. Il fenomeno della sosta diurna in mare aperto da parte di anatidi svernanti lungo le coste del Friuli Venezia Giulia. Atti I Convegno Naz. Biologi della Selvaggina. *Suppl. Ric. Biol. Selv.* 14: 89-97.
- PASCUAL C., SANCHEZ A., SANCHEZ A., VARGAS – ALBORES F., LEMOULLAC G., ROSAS C., 2003. Haemolymph metabolic variables and immune response in *Litopenaeus setiferus* adults male: the effect of an extreme temperature. *Aquaculture*, 218: 637-650.
- PAULUS K.B. & TIEDEMANN R., 2003. Ten polymorphic autosomal microsatellite loci for Eider duck *Somateria mollissima* and their cross-species applicability among waterfowl species (Anatidae). *Mol. Ecol. Notes* 3: 250 – 252.
- PEARCE J.M., TALBOT S.L., PIERSON B.J., PETERSEN M.R., SCRIBNER K.T., DICKSON D.L., MOSBECH A., 2004. Lack of spatial genetic structure among nesting and wintering. *King Eiders* 106: 229 – 240.
- PEARLSTINE E.V., 2004. Variation in mitochondrial DNA of four species of migratory raptors. *J Raptor Res* 38: 250 – 255.
- PEDDLE, D.R., 1993. An empirical comparison of Evidential reasoning, Linear Discriminant Analysis and Maximum Likelihood Algorithms for Alpine land-cover classification, *Canadian Journal of Remote Sensing*, 19: 31-44.
- PENNYCUICK C. J., 1998. Computer Simulation of Fat and Muscle Burn in Long-distance Bird Migration. *J. theor. Biol.*, 191: 47-61.
- PERCO F. & SIMONETTI G., 1999. Valle Cavanata, Foci dello Stella, Valle Canal Novo. Giunti Gruppo Editoriale, Firenze.
- PERCO F. & UTMAR P., 1989. Il censimento degli acquatici svernanti nelle principali zone umide del Friuli-Venezia Giulia fino al 1987. *Fauna* (1989) 1: 4-31.
- PERCO F. & UTMAR P., 1992. Fischione. In: Brichetti P., De Franceschi P., Baccetti N.. Aves I. Gaviidae-Phasianidae. Fauna d'Italia. Ed. Calderini. Bologna.
- PERCO F. & UTMAR P., 1997. Il censimento degli Anatidi e della Folaga svernanti nel Friuli-Venezia Giulia (1988-1996). *Fauna* 4: 23-36.
- PÉRÈS J.M., 1967. The Mediterranean Benthos. *Oceanogr. Mar. Biol. Ann. Rev.*, 5: 449-533.
- PÉRÈS J.M. & PICARD J., 1964. Nouveau Manuel de Bionomie Benthique de la Mer Mediteranee. *Rec. Trav. St. Mar. End.*, 31(47): 137.
- PEREZ-TRIS J., BENSCH S., CARBONELL R., HELBIG A.J., TELLERIA J.L., 2004. Historical diversification of migration patterns in a passerine bird. *Evolution* 58: 1819 – 1832.
- PERGENT G., MARTINI C., BOUDOURESQUE C.F., 1995. Utilisation de l'herbier à *Posidonia oceanica* comme indicateur biologique du milieu littoral en Méditerranée: état de connaissance. *Mesogee*, 54: 3-27.

- PIELOU E.C., 1966. The measurement of diversity in different types of biological collections. *J. Theor. Biol.* 13: 131-144.
- PIELOU E.C., 1969. Introduction to Mathematical Ecology. John Wiley, Sons Inc, 292 pp.
- PIERTNEY S.B., MACCOLL A.D.C., BACON P.J., RACEY P.A., LAMBIN X., DALLAS J.F., 2000. Matrilineal genetic structure and female-mediated gene flow in red grouse (*Lagopus lagopus scoticus*): an analysis using mitochondrial DNA. *Evolution* 54: 279 – 289.
- PITRA C., D'ALOIA M.-A., LIECKFELDT D., COMBREAU O., 2004. Genetic variation across the current range of the Asian houbara bustard (*Chlamydotis undulata macqueenii*). *Conserv Genet* 5: 205 – 215.
- PRATER A., MARCHANT J.J.H., VUORINEN J., 1977. Guide to the identification and ageing of Holarctic waders. British Trust for Ornithol., Field Guide #17, Tring, United Kingdom.
- PRIMMER C.R., MOLLER A.P., ELLEGREN H., 1996. A wide-range survey of cross-species microsatellite amplification in birds. *Molecular Ecology* 5, 365 – 378.
- PRUETT C.L. & WINKER K., 2005. Biological impacts of climatic change on a Beringian endemic: cryptic refugia in the establishment and differentiation of the rock sandpiper (*Calidris ptilocnemis*). *Climatic Change* 68: 219 – 240.
- RADZIEJEWSKA T. & CHABIOR M., 2004. Climatic and hydrological controls over the zoobenthos in a southern Baltic coastal lagoon. *Hydrobiologia*, 514: 171-181.
- REDFERN C.P.F., TOPP V. J., JONES P., 2004. Fat and pectoral muscle in migrating Sedge Warblers *Acrocephalus schoenobaenus*. *Ringing, Migration* 22: 24-34.
- REFISH M.M., NIGEL A.C., LANGSTON R.H.W., GREENWOOD J.J.D., 1996. A guide to the provision of refuges for waders: an analysis of 30 years of ringing data from the Wash, England. *Journal of Applied Ecology*, 33: 673-687.
- REGIONE AUTONOMA FRIULI VENEZIA GIULIA, 2008. Biodiversità. <http://www.regione.fvg.it/rafv/territorioambiente/areaArgomento.act?dir=/rafv/cms/RAFVG/AT9/ARG5/> (Query date: 27/08/2008).
- RHYMER J.M., FAIN M.G., AUSTIN J.E., JOHNSON D.H., KRAJEWSKI C., 2001. Mitochondrial phylogeography, subspecific taxonomy and conservation genetics of sandhill cranes (*Grus canadensis*; Aves: Gruidae). *Conserv Genet* 2: 203 – 218.
- RICHARDSON D.M., KAMINSKI R.M., 1992. Diet restriction, diet quality, and prebasic moult in female mallards. *J. Wildl. Manage* 56: 531-539.
- RICKLEFS R.E., 1996. Morphometry of the digestive tracts of some passerine birds. *Condor* 98, pp 279-292.
- RIEDL R., 1991. Fauna e flora del Mediterraneo, Franco Muzzio Editore.
- ROBBINS C.T., 1983. Wildlife feeding and nutrition. Academic Press, New York.
- ROGERS D.I., PIERSMA T., HASSELL C.J., 2006. Roost availability may constrain shorebird distribution: Exploring the energetic costs of roosting and disturbance around a tropical bay. *Biol. Cons.*, 133: 225-235.
- ROSE, P.M. AND SCOTT D.A. 1994. Waterfowl Population Estimates. IWRB publ. 29.
- ROWE C.L., 2002. Differences in maintenance energy expenditure by two estuarine shrimp (*Palaemonetes pugio* and *P. vulgaris*) that may permit partitioning of habitats by salinity. *Comparative Biochemistry and Physiology*, 132A: 341–351.
- RSI, 2003. *ENVI User's Guide*. RSI, Research Systems Inc. Boulder, CO. pp. 1084.
- RUOKONEN M., 2001. Phylogeography and conservation genetics of the lesser white-fronted goose (*Anser erythropus*). *Acta Universitatis Ouluensis Series a Scientiae Rerum Naturalium*. 360: 1 – 54.

- SAMBROOK J., FRITSCH E.F., MANIATIS T., 1989. Molecular cloning-a laboratory manual, 2nd edn. Cold Spring Harbor Laboratory Press, New York.
- SCHIEMANN R., 1971. Energetische Futterbewertung und Energienormen. VEB/DLV Berlin.
- SCHULZ R., STOCK M., 1993. Kentish Plovers and tourists: competitors on sandy coasts. *Wader Study Group Bull.*, 68: 83-91.
- SCRIBNER K.T., MALECKI R.A. BATT B.D.J., INMAN R.L., LIBANTS S., PRINCE H.H., 2003. Identification of source population for Greenland Canada Geese: genetic assessment of a recent colonization. *Condor* 105: 771 – 782.
- SEAMAN D.A., GUGLIELMO C.G., WILLIAMS T.D., 2006. Effects of physiological state, mass change and diet on plasma metabolite profiles in the western sandpiper (*Calidris mauri*). *The Journal of Experimental Biology* 208, pp.761-769.
- SERRA L., BACCETTI N., CHERUBINI G., ZENATELLO M., 1998. Migration and moult of Dunlin *Calidris alpina* wintering in the central Mediterranean. *Bird Study*, 45: 205-218.
- SERRA L., MAGNANI A., DALL'ANTONIA P., BACCETTI N., 1997. Risultati dei censimenti degli uccelli acquatici svernanti in Italia, 1991-1995. *Biol. Cons. Fauna*, 101: 240 pp.
- SFRISO A. & GHETTI P.F., 1998. Seasonal variation in biomass, morphometric parameters and production of seagrass in the lagoon of Venice. *Aquat. Bot.*, 61: 207-223.
- SHANNON C. E. & WEAVER W., 1949. The mathematical theory of communication. Urbana, Univ. Illinois Press: 117.
- SHANNON C.E. & WEAVER W. 1963. Mathematical theory of communication. University of Illinois Press, Urbana.
- SHEPHERD P.C.F., 2001. Space use, habitat preferences, and time activity budgets of non breeding dunlin (*Calidris alpina pacifica*) in the Frase River delta, B. C. Thesis Doctor of Philosophy Degree, Simon Fraser University, National Library of Canada.
- SHEPPARD J.K., LAWLER R. I., MARSH H., 2006. Seagrass as pasture for seacows: Landscape-level dugong habitat evaluation. *Estuarine, Coastal and Shelf Science*. Vol. 71, Issues 1-2 pp 117-132.
- SMART M. (ed.), 1974. International Conference on Conservation of Wetlands and Wildfowl, Heiligenhafen, FRG 2-6 december 1974. Proceedings. Int. Waterfowl Research Bureau (Slimbridge GB).
- SMITH H.G., WENNERBERG L., VON SCHANTZ T., 1996. Adoption or infanticide: options of replacement males in the European starling. *Behav. Ecol. Sociobiol* 38: 91 – 197.
- SMITH K.W., REED J.M., TREVIS B.E., 1992. Habitat use and site fidelity of Green Sandpipers *Tringa ochropus* wintering in southern England. *Bird Study*, 39: 155-164.
- SNOW D.W., PERRINS C.M., 1998. The Handbook of the Birds of Europe, the Middle East, and North Africa. Vol 1 non-passerines (concise edition). Oxford University Press. Oxford.
- SØNDERGAARD M., BRUUN L., LAURIDSEN T., JEPPESEN E., MADSEN T.V., 1996. The impact of grazing waterfowl on submerged macrophytes: In situ experiments in a shallow eutrophic lake. *Aquatic Botany*, 53(1): 73-84.
- SPAARGAREN D.H. & HAEFNER P.A., 1987. The effect of environmental osmotic conditions on blood and tissue glucose levels in the brown shrimp, *Crangon crangon* (L.). *Comparative Biochemistry and Physiology*, 87A, 1045–1050.
- STAI S.M. & HUGHES C.R., 2003. Characterization of microsatellite loci in wild and domestic Muscovy ducks (*Cairina moschata*). *Anim. Genet.* 34 387 – 389.

- STRAVISI F., 1988. Caratteristiche oceanografiche del Golfo di Trieste, Parco Marino di Miramare. *Hydrores*, anno V, 6: 39-45.
- SWENGEL S.R., ELLIS D.H., GEE G.F., MIRANDE C.M., 1996. Special techniques, C: Sex determination. In: *Cranes: Their Biology, Husbandry, and Conservation*; Eds.; National Biological Service/International Crane Foundation: United States of America, pp. 223 – 231.
- TAKEKAWA, J., WARNOCK N., MARTINELLI G., MILES A., TSAO D., 2002. Waterbirds use of Bayland Wetlands in the San Francisco Bay Estuary: movements of long billed dowitchers during the winter. *Waterbirds*, 25: 93-105.
- TAMISIER A., 1978a. Réalités et signification de l'organisation sociale chez les canards en période hivernale. *Bull. Soc. Fr. Etude Comport. Anim.*, 2: 11–24.
- TAMISIER A., 1978b. The functional units of wintering ducks: a spatial integration of their comfort and feeding requirements. *Verhandlungen der Ornithologischen Gesellschaft in Bayern* 23: 229-238.
- TAMISIER A. & TAMISIER M.C., 1981. L'existence d'unités fonctionnelles démontrée chez les sarcelles d'hiver en Camargue par la biotéléométrie. *Revue d'Ecologie (Terre et Vie)* 35: 563-579.
- TAMISIER A., BECHET, A., JARRY, G., LEMAHO, Y., 2003. Effets du dérangement par la chasse sur les osieaux d'eau. *Revue d'Ecologie (Terre, Vie)* 58: 435-449.
- TAN C.H. & CHOONG K.Y., 1981. Effect of hyperosmotic stress on hemolymph proteine, muscle ninhydrine – positive substances and free amino acids in *Macrobrachium rosenbergii* (De Man). *Comparative Biochemistry and Physiology*, 70 A: 485–489.
- TAYLOR, A.C. & SPICER, J.I., 1987. Metabolic response of the prawns *Palaemon elegans* and *P. serratus* (Crustacea: Decapoda) to acute hypoxia and anoxia. *Marine Biology*, 95: 521 – 530.
- TIEDEMANN R., 1999. Seasonal changes in the breeding origin of migrating Dunlins (*Calidris alpina*) as revealed by mitochondrial DNA sequencing. *J Ornithol.* 140: 319 – 323.
- TORRES G., GIMENEZ L., ANGER K., 2002. Effects of reduced salinity on the biochemical composition (lipid, protein) of zoea decapod crustacean larvae. *Journal of Experimental Marine Biology and Ecology*, 277: 43-60.
- TOTZKE U. & BAIRLEIN F., 1998. The body mass cycle of the migratory garden warbler (*Sylvia borin*) is associated with changes of basal plasma metabolite levels. *Comparative Biochemistry and Physiology Part A* 121:127-133.
- TRIPLET P., SOURNIA A., JOYEUX E., LE DREAN-QUENEC'H DU S., 2003. Activités humaines et dérangement: l'exemple des osieaux d'eau. *Alauda* 71: 305-316.
- TUCKER M. & EVANS M.I., 1997. Habitat for birds in Europe: a conservation strategy for the wider environment. Cambridge, U.K.: BirdLife International (BirdLife Conservation Series no 6).
- TURCEK F.J., 1956. Zur Frage der Dominanz in Vogelpopulationen. *Waldhygiene* 8: 249-257.
- TURPIE J. K. & HOCKEY P.A.R., 1996. Foraging ecology and seasonal energy budgets of estuarine Grey Plovers *Pluvialis squatarola* and Whimbrels *Numenius phaeopus* at the southern tip of Africa. *Ardea*, 84: 57-74.
- URQUHART M., ARMOUR J., DUNCAN J.L., DUNN A.M., JENNINGS F.W., 1998. *Parassitologia Veterinaria*. UTET, Torino.
- VANGELUWE D., BEUDELS M.O., LAMANI F., 1994. Evaluation de l'état de conservation des zones humides du littoral albanais et appréciation de leurs potentialité pour le Courlis à bec grêle *Numenius tenuirostris*, in (1994) : Préparation d'un plan de sauvetage pour *Numenius tenuirostris*, Rapport à la Commission des Communautés européennes. Relazione inedita.

- VERSLYCKE T. & JANSSEN C.R., 2002. Effects of a changing abiotic environment on the energy metabolism in the estuarine mysid shrimp *Neomysis integer* (Crustacea: Mysidacea). *Journal of Experimental Marine Biology and Ecology*, 279: 61–72.
- VON KÄNEL A., 1981. Winter feeding ecology of Wigeon (*Anas penelope*) at the Ouse Washes, England. *Ibis*, 123(3):438–449.
- WAITS J.L., AVERY M.L., TOBIN M.E., LEBERG P.L., 2003. Low mitochondrial DNA variation in double-crested cormorants in Eastern North America. *Water birds* 26: 196 – 200.
- WALDENSTROM J., BENSCH S., KIBOI S., HASSELQUIST D., OTTOSSON U., 2002. Cross-species infection of blood parasites between resident and migratory songbirds in Africa. *Mol Ecol* 11: 1545 – 1554.
- WARNOCK S.E., TAKEKAWA J.Y., 1996. Wintering site fidelity and movement patterns of Western Sandpipers *Calidris mauri* in the San Francisco Bay estuary. *Ibis*, 138: 160-167.
- WELSH J. AND MC CLEALLAND M., 1990. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acid Research* 18: 7213 – 7218.
- WENINK P.W. & BAKER A.J., 1996. Mitochondrial DNA lineages in composite flocks of migratory and wintering dunlins (*Calidris alpina*). *Auk* 113 744 – 756.
- WENNERBERG L., 2001. Breeding origin and migration pattern of dunlin (*Calidris alpina*) revealed by mitochondrial DNA analysis. *Mol Ecol* 10 1111 – 1120.
- WENNERBERG L., HOLMGREN N.M.A., JONSSON P.-E., VON SCHANTZ T., 1999. Genetic and morphological variation in Dunlin *Calidris alpina* breeding in the Palearctic tundra. *Ibis* 141: 391 – 398.
- WENNERBERG L., KLAASSEN M., LINDSTROM A., 2002. Geographical variation and population structure in the White-rumped Sandpiper *Calidris fuscicollis* as shown by morphology, mitochondrial DNA and carbon isotope ratios. *Oecologia* 131: 380 – 390.
- WIENS J.A., 1975. Avian communities, energetics and functions in coniferous forests habitats. Proc. Symp. Manag. Forest Range Habitats Nongame Birds, Tucson. USDA Forest Service: 146-182.
- WIKIPEDIA, 2008. Adriatico. http://it.wikipedia.org/wiki/Mar_Adriatico (Query date: 21/08/2008).
- WILLIAMS T.D., GUGLIELMO C.G., EGELER O., MARTYNIUK C.J., 1999. Plasma lipid metabolites provide information on mass change over several days in captive Western Sandpipers. *The Auk* 116 pp.994-1000.
- WINK M., 2006. Use of DNA markers to study bird migration. *J Ornithol* 147: 234 – 244.
- WINK M., SAUER-GÜRTH H., GWINNER E., 2002. A molecular phylogeny of stonechats and related turdids inferred from mitochondrial DNA sequences and genomic fingerprinting by ISSR-PCR. *Brit Birds* 95: 349 – 355.
- WORLD COMMISSION ON PROTECTED AREAS, 2005. *World database on protected areas*. Available: <http://sea.unep-wcmc.org/wdbpa/> (Query date: 28/08/08).
- ZAR JH, 1999. *Biostatistical analysis*. 4th edn. Prentice-Hall, New Jersey.
- ZWARTS L. & ESSELINK P., 1989. Versatility of male curlews *Numenius arquata* preying upon *Nereis diversicolor*: deploying contrasting capture modes dependent on prey availability. *Marine Ecology Progress Series*, 56: 255-269.